

For the treatment of adult men with Peyronie's disease who have a "plaque" that can be felt and a curve in their penis greater than 30 degrees when treatment is started

What to expect during treatment WITH XIAFLEX[®]

IMPORTANT SAFETY INFORMATION FOR XIAFLEX[®]

XIAFLEX[®] can cause serious side effects, including penile fracture (corporal rupture) or other serious injury to the penis. Receiving an injection of XIAFLEX[®] may cause damage to the tubes in your penis called the corpora. After treatment with XIAFLEX[®], one of these tubes may break during an erection. This is called a corporal rupture or penile fracture. This could require surgery to fix the damaged area. Damage to your penis might not get better after a corporal rupture.

Please see additional Important Safety Information throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

XIAFLEX[®]
collagenase clostridium histolyticum

XIAFLEX® is a prescription medicine used to treat adult men with Peyronie's disease who have a "plaque" that can be felt and a curve in their penis greater than 30 degrees when treatment is started.



Actor portrayals.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

Do not receive XIAFLEX® if you:

- have been told by your healthcare provider that the Peyronie's plaque to be treated involves the "tube" that your urine passes through (urethra)
- are allergic to collagenase clostridium histolyticum or any of the ingredients in XIAFLEX®, or to any other collagenase product. See the end of the Medication Guide for a complete list of ingredients in XIAFLEX®

Please see the Important Safety Information on pages 11-13 and throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

Welcome to treatment with XIAFLEX®

Now that you've been prescribed XIAFLEX®, you and your urologist will become partners as you begin treatment. Here's what you should be prepared to do before treatment starts:



ANSWER THE PHONE IF THE PHARMACY CALLS

- You may need to verify coverage for treatment with XIAFLEX®
- If you miss the call, simply **call US Bioservices at 1-888-518-7246** to verify your benefits



SCHEDULE YOUR APPOINTMENTS

- Be sure to schedule all your appointments with your urologist
- You can use the Appointment Card on the last page to keep track of each visit

Starting your treatment

You and your urologist will participate in 2 equally important parts of your treatment.

PART 1: WORKING WITH YOUR HCP



Visit 1: First injection of XIAFLEX®

- Your healthcare professional (HCP) will inject XIAFLEX® into the plaque that is causing your penis to curve
- XIAFLEX® is given as part of a treatment cycle. In each treatment cycle, you will receive an injection of XIAFLEX®, followed by a second injection 1 to 3 days later



Visit 2: Second injection of XIAFLEX®

- Occurs 1 to 3 days after initial injection



Visit 3: HCP penile modeling

- 1 to 3 days after the second injection of XIAFLEX®, you will need to return to your HCP's office for a manual procedure called penile modeling that will stretch and help straighten your penis



Actor portrayals.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

XIAFLEX® can cause serious side effects, including:

1. **Penile fracture (corporal rupture) or other serious injury to the penis.** Receiving an injection of XIAFLEX® may cause damage to the tubes in your penis called the corpora. After treatment with XIAFLEX®, one of these tubes may break during an erection. This is called a corporal rupture or penile fracture. This could require surgery to fix the damaged area. Damage to your penis might not get better after a corporal rupture.

XIAFLEX® should be injected directly into the plaque that is causing your erect penis to curve by a urologist who is trained and experienced in treating adult men with Peyronie's disease.

Please see the Important Safety Information on pages 11-13 and throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

XIAFLEX®
collagenase clostridium histolyticum

Starting your treatment (cont)

PART 2: YOUR ROLE AT HOME



Daily penile modeling

- After each treatment cycle, you will need to take a few minutes each day to perform at-home stretching and straightening activities on your penis for about 6 weeks
- To find out how to perform these activities, see the daily penile modeling activities section on pages 8 and 9 of this brochure



Sexual activity

For each treatment cycle, do not have sex or engage in any other sexual activity:

- Between the first and second injections
- For at least 4 weeks after the second injection and after any pain and swelling has gone away

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

- After treatment with XIAFLEX®, blood vessels in your penis may also break, causing blood to collect under the skin (hematoma). This could require a procedure to drain the blood from under the skin

Symptoms of corporal rupture or other serious injury to your penis may include:

- a popping sound or sensation in an erect penis
- sudden loss of the ability to maintain an erection
- pain in your penis
- purple bruising and swelling of your penis
- difficulty urinating or blood in the urine



After each injection

- Your penis may be wrapped with a bandage. Your HCP will tell you when to take the bandage off
- It's common to see bruising and swelling after your injections with XIAFLEX®
- Avoid situations that may cause you to strain your stomach (abdominal) muscles, such as straining during bowel movements
- **Do not use a vacuum erection device during your treatment with XIAFLEX®**

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

Call your healthcare provider right away if you have any of the symptoms of corporal rupture or serious injury to the penis listed on the left.

Please see the Important Safety Information on pages 11-13 and throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

This completes 1 of 4 possible treatment cycles. You and your urologist will determine how many cycles are appropriate for you.

Your role during treatment

How to perform daily penile modeling activities at home

Part of your commitment to treatment means performing the 2 types of penile modeling activities (stretching and straightening) at home daily for 6 weeks at the end of each treatment cycle.

STRETCHING ACTIVITY

3x per day | when your penis is not hard (erect)



Gently stretch your penis 3 times a day. Only stretch your penis if your penis is not hard (erect).

1. With one hand, hold the tip of your penis with your fingers. With your other hand, hold the base of your penis with your fingers.
2. Gently pull your penis away from your body to its full length and hold the stretch for 30 seconds.
3. Let go of the tip of your penis and let your penis return to its normal length.

STRETCHING TIP:

- It might be helpful to set aside times during the day when you have privacy and will not be interrupted

STRAIGHTENING ACTIVITY

1x per day | when your penis is erect (unrelated to sexual activity)



Gently straighten your penis once a day. Only straighten your penis if you have an erection that happens without any sexual activity (spontaneous erection). Bending your penis should not cause any pain or discomfort.

1. With one hand, hold your erect penis.
2. With your other hand, gently bend your penis in the opposite direction of the curve.
3. Hold the penis in this more straightened position for 30 seconds, then let go.

STRAIGHTENING TIP:

- Perform this activity in the morning if you experience a spontaneous erection

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

Do not have sex or any other sexual activity between the first and second injections of a treatment cycle.

Do not have sex or have any other sexual activity for at least 4 weeks after the second injection of a treatment cycle with XIAFLEX® and after any pain and swelling have gone away.

Please see the Important Safety Information on pages 11-13 and throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

XIAFLEX®
collagenase clostridium histolyticum

Follow these helpful tips to get the most out of your treatment

BEFORE TREATMENT



Answer the phone if the pharmacy calls

- You may need to verify coverage for treatment with XIAFLEX®
- If you miss the call, simply **call US Bioservices at 1-888-518-7246** to verify your benefits



Schedule all of your appointments

Work with your HCP to schedule all of your treatments in advance

DURING TREATMENT



Keep track of appointments and your daily penile modeling

Using a calendar or app may help you stay on top of things



Perform daily penile modeling activities properly

- Your HCP will show you how to perform these activities
- For a refresher, refer to pages 8 and 9 of this brochure



Talk to your partner

You may find it helpful to discuss how you're feeling and involve them in the treatment process



Stay in touch with your urologist

Be sure to contact your urologist if you have any questions, concerns, or experience any side effects

WHAT IS XIAFLEX®?

XIAFLEX® is a prescription medicine used to treat adult men with Peyronie's disease who have a "plaque" that can be felt and a curve in their penis greater than 30 degrees when treatment is started.

It is not known if XIAFLEX® is safe and effective in children under the age of 18.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX®

Do not receive XIAFLEX® if you:

- have been told by your healthcare provider that the Peyronie's plaque to be treated involves the "tube" that your urine passes through (urethra)
- are allergic to collagenase clostridium histolyticum or any of the ingredients in XIAFLEX®, or to any other collagenase product. See the end of the Medication Guide for a complete list of ingredients in XIAFLEX®

XIAFLEX® can cause serious side effects, including:

1. **Penile fracture (corporal rupture) or other serious injury to the penis.** Receiving an injection of XIAFLEX® may cause damage to the tubes in your penis called the corpora. After treatment with XIAFLEX®, one of these tubes may break during an erection. This is called a corporal rupture or penile fracture. This could require surgery to fix the damaged area. Damage to your penis might not get better after a corporal rupture.
 - After treatment with XIAFLEX®, blood vessels in your penis may also break, causing blood to collect under the skin (hematoma). This could require a procedure to drain the blood from under the skin

(continued on next page)

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

XIAFLEX®
collagenase clostridium histolyticum

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

Symptoms of corporal rupture or other serious injury to your penis may include:

- a popping sound or sensation in an erect penis
- sudden loss of the ability to maintain an erection
- pain in your penis
- purple bruising and swelling of your penis
- difficulty urinating or blood in the urine

Call your healthcare provider right away if you have any of the symptoms of corporal rupture or serious injury to the penis listed above.

Do not have sex or any other sexual activity between the first and second injections of a treatment cycle.

Do not have sex or have any other sexual activity for at least 4 weeks after the second injection of a treatment cycle with XIAFLEX® and after any pain and swelling have gone away.

XIAFLEX® for the treatment of Peyronie's disease is only available through a restricted program called the XIAFLEX® Risk Evaluation and Mitigation Strategy (REMS) Program. For more information about the XIAFLEX® REMS Program, go to www.XIAFLEXREMS.com or call 1-877-313-1235.

2. Hypersensitivity reactions, including anaphylaxis. Severe allergic reactions can happen in people who receive XIAFLEX®, because it contains foreign proteins.

Call your healthcare provider right away if you have any of these symptoms of an allergic reaction after an injection of XIAFLEX®:

- hives
- swollen face
- breathing trouble
- chest pain
- low blood pressure
- dizziness or fainting

Before receiving XIAFLEX®, tell your healthcare provider if you have had an allergic reaction to a previous XIAFLEX® injection, or have a bleeding problem or any other medical conditions. Tell your healthcare provider about all the

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX® (cont)

medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Using XIAFLEX® with certain other medicines can cause serious side effects. Especially tell your healthcare provider if you take medicines to thin your blood (anticoagulants). If you are told to stop taking a blood thinner before your XIAFLEX® injection, your healthcare provider should tell you when to restart the blood thinner. Ask your healthcare provider or pharmacist for a list of these medicines if you are unsure.

What should I avoid while receiving XIAFLEX®?

Avoid situations that may cause you to strain your stomach (abdominal) muscles, such as straining during bowel movements.

Do not use a vacuum erection device during your treatment with XIAFLEX®.

XIAFLEX® can cause serious side effects, including increased chance of bleeding. Bleeding or bruising at the injection site can happen in people who receive XIAFLEX®. Talk to your healthcare provider if you have a problem with your blood clotting. XIAFLEX® may not be right for you.

The most common side effects with XIAFLEX® for the treatment of Peyronie's disease include:

- a small collection of blood under the skin at the injection site (hematoma)
- swelling at the injection site or along your penis
- pain or tenderness at the injection site, along your penis and above your penis
- penis bruising
- itching of your penis or scrotum (genitals)
- painful erection
- erection problems (erectile dysfunction)
- changes in the color of the skin of your penis
- blisters at the injection site
- pain with sex
- a lump at the injection site (nodule)

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all of the possible side effects with XIAFLEX®. For more information, ask your healthcare provider or pharmacist.

XIAFLEX®
collagenase clostridium histolyticum

XIAFLEX® Copay Assistance Program

Patient eligibility

Patients qualify for assistance under this program if:

- You are receiving or have received XIAFLEX® for an approved indication and in a manner consistent with the instructions for administration of XIAFLEX®
- You are uninsured or have insurance that is not provided by Medicare, Medicare Prescription Drug Benefit plans, Medicare Advantage, Veterans Affairs (VA), Medicaid, or similar federal or state programs, and this program is not otherwise prohibited by law
- You are 18 years of age or older
- You have paid or are obligated to pay out-of-pocket costs for a prescription of XIAFLEX®

No other purchase is necessary to receive this offer.

For more information, call 1-800-743-2382.

Restrictions, Terms and Conditions

1. By accepting this offer, you agree to report the value received under this offer to any health insurer or other third party paying for any part of your XIAFLEX® prescription if you are required to do so by benefit terms, contract, or law.
2. This offer is not valid for prescriptions reimbursed in whole or in part by Medicare, Medicare Prescription Drug Benefit plans, Medicare Advantage, VA, Medicaid, or similar federal or state programs, or where otherwise prohibited by law.
3. By accepting this offer, you agree that Endo Pharmaceuticals Inc. or those working on its behalf may contact your doctor to verify information about treatment that is relevant to verifying your eligibility for this offer.
4. This offer is only valid for doses of XIAFLEX® administered in the US.
5. This offer is valid for the out-of-pocket cost for a dose of XIAFLEX® only. Offer is not valid for any other products or other out-of-pocket costs (for example, office visit charges, office visit copays, or injection/administration costs), even if those costs are associated with the administration of a dose of XIAFLEX®.
6. This offer is valid only if you have not used this program within the last 30 days.
7. The selling, purchasing, trading, or counterfeiting of this offer is prohibited.
8. Endo Pharmaceuticals Inc. reserves the right to rescind, revoke, or amend this offer without notice.
9. By participating, you understand and agree to comply with the terms and conditions of this offer as set forth above.

Please see the Important Safety Information on pages 11-13 and throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

Learn more
about XIAFLEX®
at xiaflex.com



Actor portrayals.

If eligible,
you'll pay **\$0 OUT OF
POCKET**

Most eligible patients with Commercial insurance plans paid \$0 copay* for XIAFLEX®.

To find out if you qualify,
call 1-800-743-2382

*Based on analysis of over 4000 claims filed
January 2015 through October 2015.

For more information,
visit xiaflex.com or
call 877-XIAFLEX
(877-942-3539)

XIAFLEX[®]
collagenase clostridium histolyticum

For the treatment of adult men with Peyronie's disease who have a "plaque" that can be felt and a curve in their penis greater than 30 degrees when treatment is started

IMPORTANT SAFETY INFORMATION FOR XIAFLEX[®]

XIAFLEX[®] can cause serious side effects, including penile fracture (corporal rupture) or other serious injury to the penis. Receiving an injection of XIAFLEX[®] may cause damage to the tubes in your penis called the corpora. After treatment with XIAFLEX[®], one of these tubes may break during an erection. This is called a corporal rupture or penile fracture. This could require surgery to fix the damaged area. Damage to your penis might not get better after a corporal rupture.

Please see additional Important Safety Information throughout this brochure.

Please [click here](#) for full Prescribing Information, including Boxed Warning and Medication Guide.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.

 **endo**
pharmaceuticals
an endo international company

Rx Only

XIAFLEX[®] is a registered trademark of Endo International plc or one of its affiliates.
© 2019 Endo Pharmaceuticals Inc. All rights reserved. Malvern, PA 19355
XP-05592/August 2019 www.xiaflex.com 1-800-462-ENDO (3636)

Keep track of visits with your healthcare professional (HCP)

APPOINTMENT CARD

CYCLE 1

1st Injection: ___ / ___ / ___

2nd Injection: ___ / ___ / ___

HCP Modeling: ___ / ___ / ___

CYCLE 2

1st Injection: ___ / ___ / ___

2nd Injection: ___ / ___ / ___

HCP Modeling: ___ / ___ / ___

CYCLE 3

1st Injection: ___ / ___ / ___

2nd Injection: ___ / ___ / ___

HCP Modeling: ___ / ___ / ___

CYCLE 4

1st Injection: ___ / ___ / ___

2nd Injection: ___ / ___ / ___

HCP Modeling: ___ / ___ / ___

You and your urologist will determine how many cycles are appropriate for you.